

PHARMACY POLICY STATEMENT

Indiana Medicaid

		transplant (HSCT), od
DRUG NAME	Rezurock (belumosudil)	at least half of patier
BLLING CODE	Must use alid NDC	regarding optimal 2
BNEFIT TIPE	Pharmacy	regarding optimal 2

nd line therapy but describe ariety of options.

Rezurockis the first rho-associated, coiled-coil kinase 2 (ROCK) inhibitor. The ROCK pathway modulates inflammatory response and fibrotic processes. ROCK inhibition is thought to both restore immune homeostasis and reduce fibrotic processes, which makes Rezurociqe from other pharmacologic treatment options. Approxi was based on the phase 2 ROCK at study.

Rezurock (belumosudilyill be considered for coerage when the following criteria are met:

Chronic Graft-Versus-Host Disease (cGVHD)

For initial authorization:

- 1. Member is at least 12 yearsof age; AND
- 2. Medication must be prescribed by or in consultation with an splant or hematology/oncology specialist; AND
- 3 Member has adiagnosis of GVHD with persistent manifestation ASND
- 4. Member has failed at least 2 prior lines of systemic therapy, i.e., systemic cot

For reauthorization:

1. Chart notes must show improvement of signs and symptoms of disease in at least 1 organ/site, without progression in any other organ/site.

If all the above requirements are met, the medication will be approved for an additional 12 months.

IN-MED-P-366647 OMPP Approved .383